

Up to **four** people can work on an exercise together. But each of you should be able to explain the solutions to the TA (Bremser). Write your names **and** the name of your group (time, TA) on the sheets. Staple them together

## Assignment 11

Deadline: February 9, 2004

Solve at least two of the following three exercises.

### Exercise 1

In this exercise we discuss how to convert a minimization problem into a decision problem. Assume that the cost function has nonnegative integer values and that we know an upper bound  $\hat{f} \geq f^*$  for the value of an optimal solution. Further assume that we have a program  $\mathcal{P} : \mathbb{N} \rightarrow \mathcal{L} \cup \{\perp\}$  so that  $\mathcal{P}(b)$  outputs a feasible solution  $\mathbf{x} \in \mathcal{L}$  with  $f(\mathbf{x}) \leq b$  if such a solution exists and signals failure  $\mathcal{P}(b) = \perp$  otherwise.

- (a) Explain how to find an optimal solution using  $O(\log \hat{f})$  calls of  $\mathcal{P}$ .
- (b) Refine the procedure from part (a) so that it works using  $O(\log f^*)$  executions of  $\mathcal{P}$ , even if we do not know an upper bound of  $f^*$ .

### Exercise 2

Suppose we wish to invest 14,000 euros for one year. We have identified four investment opportunities. Investment 1 requires an investment of 5,000 euros and gives a profit of 800 euros; investment 2 requires 7,000 euros and gives a profit of 1,100 euros; investment 3 requires 4,000 euros and gives a profit of 600 euros; and investment 4 requires 3,000 euros and gives a profit of 400 euros. Into which investments should we place our money so as to maximize our total profit? Model the problem as a knapsack problem. (Hint: For convenience, drop extra zeros.)

### Exercise 3

Consider the following algorithm for the traveling salesman problem for undirected graphs with triangle inequality (see the lecture slides for a definition):

- Let  $T$  denote the edges of a minimum spanning tree of  $G$ .
- Now consider the directed graph  $G' = (V, T')$  where for each edge  $\{u, v\} \in T$ ,  $T'$  contains the two directed edges  $(u, v)$  and  $(v, u)$ . Note that each node in  $(V, T')$  has the same in-degree as out-degree.
- Find an Euler tour  $C'$  in  $G'$ .
- Convert  $C'$  into a simple cycle visiting all nodes by introducing *shortcuts*, i.e., repeatedly apply the following shortcutting step: If there are subsequent edges  $(u, v)$  and  $(v, w)$  in  $C'$  such that  $v$  is also visited before, replace these two edges by the edge  $(u, w)$ .

Now we ask you to show that this algorithm yields a two-approximation of the optimal TSP tour. (Hint: you might need to show that the weight of  $T$  is no more than the weight of an optimal traveling salesman tour.) Do not forget to argue the above algorithm works in polynomial time.