2.10 Refutational Completeness of Resolution

How to show refutational completeness of propositional resolution:

- We have to show: $N \models \bot \Rightarrow N \vdash_{Res} \bot$, or equivalently: If $N \nvdash_{Res} \bot$, then N has a model.
- Idea: Suppose that we have computed sufficiently many inferences (and not derived \perp).
- Now order the clauses in N according to some appropriate ordering, inspect the clauses in ascending order, and construct a series of Herbrand interpretations.
- \bullet The limit interpretation can be shown to be a model of N.

Clause Orderings

- 1. We assume that ≻ is any fixed ordering on ground atoms that is *total* and *well-founded*. (There exist many such orderings, e.g., the lenght-based ordering on atoms when these are viewed as words over a suitable alphabet.)
- 2. Extend \succ to an ordering \succ_L on ground literals:

$$[\neg]A \succ_L [\neg]B$$
, if $A \succ_B \neg A \succ_L A$

3. Extend \succ_L to an ordering \succ_C on ground clauses: $\succ_C = (\succ_L)_{\text{mul}}$, the multi-set extension of \succ_L .

Notation: \succ also for \succ_L and \succ_C .

Example

Suppose $A_5 > A_4 > A_3 > A_2 > A_1 > A_0$. Then:

$$A_0 \lor A_1$$

$$\prec A_1 \lor A_2$$

$$\prec \neg A_1 \lor A_2$$

$$\prec \neg A_1 \lor A_4 \lor A_3$$

$$\prec \neg A_1 \lor \neg A_4 \lor A_3$$

$$\prec \neg A_5 \lor A_5$$


Properties of the Clause Ordering

Proposition 2.20:

- 1. The orderings on literals and clauses are total and well-founded.
- 2. Let C and D be clauses with $A = \max(C)$, $B = \max(D)$, where $\max(C)$ denotes the maximal atom in C.
 - (i) If $A \succ B$ then $C \succ D$.
 - (ii) If A = B, A occurs negatively in C but only positively in D, then $C \succ D$.

Stratified Structure of Clause Sets

Let A > B. Clause sets are then stratified in this form:



Closure of Clause Sets under Res

 $Res(N) = \{C \mid C \text{ is concl. of a rule in } Res \text{ w/ premises in } N\}$ $Res^0(N) = N$

$$Res^{n+1}(N) = Res(Res^n(N)) \cup Res^n(N)$$
, for $n \ge 0$
 $Res^*(N) = \bigcup_{n \ge 0} Res^n(N)$

N is called saturated (wrt. resolution), if $Res(N) \subseteq N$.

Proposition 2.21:

- (i) $Res^*(N)$ is saturated.
- (ii) Res is refutationally complete, iff for each set N of ground clauses:

$$N \models \bot \Leftrightarrow \bot \in Res^*(N)$$

Construction of Interpretations

Given: set N of ground clauses, atom ordering \succ . Wanted: Herbrand interpretation I such that

- "many" clauses from N are valid in I;
- $I \models N$, if N is saturated and $\bot \notin N$.

Construction according to \succ , starting with the minimal clause.

Example

Let $A_5 \succ A_4 \succ A_3 \succ A_2 \succ A_1 \succ A_0$ (max. literals in red)

	clauses <i>C</i>	I _C	Δ_{C}	Remarks
1	$\neg A_0$	Ø	Ø	true in I_C
2	$A_0ee A_1$	Ø	$\{A_1\}$	A_1 maximal
3	$A_1ee A_2$	$\{A_1\}$	Ø	true in I_C
4	$ eg \mathcal{A}_1 ee \mathcal{A}_2$	$\{A_1\}$	$\{A_2\}$	A_2 maximal
5	$\neg A_1 \lor A_4 \lor A_3 \lor A_0$	$\{A_1,A_2\}$	$\{A_4\}$	A_4 maximal
6	$\neg A_1 \lor \neg A_4 \lor A_3$	$\{A_1, A_2, A_4\}$	Ø	A_3 not maximal;
				min. counter-ex.
7	$ eg \mathcal{A}_1 ee \mathcal{A}_5$	$\{A_1, A_2, A_4\}$	$\{A_5\}$	

 $\overline{I} = \{A_1, A_2, A_4, A_5\}$ is not a model of the clause set \Rightarrow there exists a counterexample.

Main Ideas of the Construction

- \bullet Clauses are considered in the order given by \prec .
- When considering C, one already has a partial interpretation I_C (initially $I_C = \emptyset$) available.
- If C is true in the partial interpretation I_C , nothing is done. $(\Delta_C = \emptyset)$.
- If C is false, one would like to change I_C such that C becomes true.

Main Ideas of the Construction

- Changes should, however, be monotone. One never deletes anything from I_C and the truth value of clauses smaller than C should be maintained the way it was in I_C .
- Hence, one chooses $\Delta_C = \{A\}$ if, and only if, C is false in I_C , if A occurs positively in C (adding A will make C become true) and if this occurrence in C is strictly maximal in the ordering on literals (changing the truth value of A has no effect on smaller clauses).

Resolution Reduces Counterexamples

$$\frac{\neg A_1 \lor A_4 \lor A_3 \lor A_0 \quad \neg A_1 \lor \neg A_4 \lor A_3}{\neg A_1 \lor \neg A_1 \lor A_3 \lor A_3 \lor A_0}$$

Construction of *I* for the extended clause set:

clauses C	I _C	Δ_{C}	Remarks
$\neg A_0$	Ø	Ø	
$A_0 \lor A_1$	Ø	$\{A_1\}$	
$A_1 \lor A_2$	$\{\mathcal{A}_1\}$	Ø	
$ eg \mathcal{A}_1 ee \mathcal{A}_2$	$\{\mathcal{A}_1\}$	$\{A_2\}$	
$\neg A_1 \lor \neg A_1 \lor A_3 \lor A_3 \lor A_0$	$\{A_1,A_2\}$	Ø	A_3 occurs twice
			minimal counter-ex.
$ eg A_1 \lor A_4 \lor A_3 \lor A_0$	$\{A_1,A_2\}$	$\{A_4\}$	
$\neg A_1 \lor \neg A_4 \lor A_3$	$\{A_1, A_2, A_4\}$	Ø	counterexample
$ eg A_1 \lor A_5$	$\{A_1, A_2, A_4\}$	$\{A_5\}$	

The same I, but smaller counterexample, hence some progress was made.

Factorization Reduces Counterexamples

$$\frac{\neg A_1 \vee \neg A_1 \vee A_3 \vee A_3 \vee A_0}{\neg A_1 \vee \neg A_1 \vee A_3 \vee A_0}$$

Construction of *I* for the extended clause set:

clauses C	I _C	Δ_{C}	Remarks
$\neg A_0$	Ø	Ø	
$A_0 \lor A_1$	Ø	$\{\mathcal{A}_1\}$	
$A_1 \lor A_2$	$\{A_1\}$	Ø	
$ eg \mathcal{A}_1 ee \mathcal{A}_2$	$\{A_1\}$	$\{A_2\}$	
$ eg A_1 \lor eg A_1 \lor eg A_3 \lor eg A_0$	$\{A_1, A_2\}$	$\{A_3\}$	
$\neg A_1 \lor \neg A_1 \lor A_3 \lor A_3 \lor A_0$	$\{A_1, A_2, A_3\}$	Ø	true in I_C
$ eg A_1 \lor A_4 \lor A_3 \lor A_0$	$\{A_1, A_2, A_3\}$	Ø	
$\neg A_1 \lor \neg A_4 \lor A_3$	$\{A_1, A_2, A_3\}$	Ø	true in I_C
$\neg A_3 \lor A_5$	$\{A_1, A_2, A_3\}$	$\{A_5\}$	

The resulting $I = \{A_1, A_2, A_3, A_5\}$ is a model of the clause set.

Construction of Candidate Models Formally

Let N, \succ be given. We define sets I_C and Δ_C for all ground clauses C over the given signature inductively over \succ :

$$I_C := \bigcup_{C \succ D} \Delta_D$$

$$\Delta_C := \begin{cases} \{A\}, & \text{if } C \in N, \ C = C' \lor A, \ A \succ C', \ I_C \not\models C \\ \emptyset, & \text{otherwise} \end{cases}$$


We say that C produces A, if $\Delta_C = \{A\}$.

The candidate model for N (wrt. \succ) is given as $I_N^{\succ} := \bigcup_C \Delta_C$.

We also simply write I_N , or I, for I_N^{\succ} if \succ is either irrelevant or known from the context.

Structure of N, \succ

Let A > B; producing a new atom does not affect smaller clauses.



Some Properties of the Construction

Proposition 2.22:

- (i) $C = \neg A \lor C' \Rightarrow \text{no } D \succeq C \text{ produces } A.$
- (ii) C productive $\Rightarrow I_C \cup \Delta_C \models C$.
- (iii) Let $D' \succ D \succeq C$. Then

$$I_D \cup \Delta_D \models C \Rightarrow I_{D'} \cup \Delta_{D'} \models C$$
 and $I_N \models C$.

If, in addition, $C \in N$ or $max(D) \succ max(C)$:

$$I_D \cup \Delta_D \not\models C \Rightarrow I_{D'} \cup \Delta_{D'} \not\models C$$
 and $I_N \not\models C$.

Some Properties of the Construction

(iv) Let $D' \succ D \succ C$. Then

$$I_D \models C \Rightarrow I_{D'} \models C$$
 and $I_N \models C$.

If, in addition, $C \in N$ or $max(D) \succ max(C)$:

$$I_D \not\models C \Rightarrow I_{D'} \not\models C$$
 and $I_N \not\models C$.

(v) $D = C \vee A$ produces $A \Rightarrow I_N \not\models C$.

Model Existence Theorem

Theorem 2.23 (Bachmair & Ganzinger):

Let \succ be a clause ordering, let N be saturated wrt. Res, and suppose that $\bot \notin N$. Then $I_N^{\succ} \models N$.

Corollary 2.24:

Let N be saturated wrt. Res. Then $N \models \bot \Leftrightarrow \bot \in N$.

Model Existence Theorem

Proof of Theorem 2.23:

Suppose $\bot \notin N$, but $I_N^{\succ} \not\models N$. Let $C \in N$ minimal (in \succ) such that $I_N^{\succ} \not\models C$. Since C is false in I_N , C is not productive. As $C \neq \bot$ there exists a maximal atom A in C.

Case 1: $C = \neg A \lor C'$ (i.e., the maximal atom occurs negatively) $\Rightarrow I_N \models A$ and $I_N \not\models C'$

 \Rightarrow some $D = D' \lor A \in N$ produces A. As $\frac{D' \lor A}{D' \lor C'}$, we infer that $D' \lor C' \in N$, and $C \succ D' \lor C'$ and $I_N \not\models D' \lor C'$ \Rightarrow contradicts minimality of C.

Case 2: $C = C' \lor A \lor A$. Then $\frac{C' \lor A \lor A}{C' \lor A}$ yields a smaller counterexample $C' \lor A \in N$. \Rightarrow contradicts minimality of C.

Compactness of Propositional Logic

Theorem 2.25 (Compactness):

Let N be a set of propositional formulas. Then N is unsatisfiable, if and only if some finite subset $M \subseteq N$ is unsatisfiable.

Proof:

"⇐": trivial.

" \Rightarrow ": Let N be unsatisfiable.

 $\Rightarrow Res^*(N)$ unsatisfiable

 $\Rightarrow \bot \in Res^*(N)$ by refutational completeness of resolution

 $\Rightarrow \exists n \geq 0 : \bot \in Res^n(N)$

 $\Rightarrow \bot$ has a finite resolution proof P;

choose M as the set of assumptions in P.