

6.2 Linear Arithmetic

There are several ways of introducing linear arithmetic and in particular its syntax. I start with a syntax that already contains $-$, \leq , $<$, \geq , \neq and \mathbb{Q} . All these functions and relations are indeed expressible by first-order formulas over 0 , 1 , \approx , and $>$. For the semantics there are two approaches. Either providing axioms, i.e., closed formulas, for the above symbols and then considering all algebras satisfying the axioms, or fixing one particular algebra or a class of algebras. For this chapter I start with a rich syntax and a semantics based on a fixed algebra.

Definition 6.2.1 (LA Syntax). The syntax of LA is

$$\Sigma_{\text{LA}} = (\{\text{LA}\}, \{0, 1, +, -\} \cup \mathbb{Q}, \{\leq, <, \neq, >, \geq\})$$

where $-$ is unitary and all other symbols have the usual arities.

Terms and formulas over Σ_{LA} are built in the classical free first-order way, see Section 3.1. All first-order notions, i.e., terms, atoms, equations, literals, clauses, etc. carry over to LA formulas. The atoms and terms built over the LA signature are written in their standard infix notation, i.e., I write $3 + 5$ instead of $+(3, 5)$. Note that the signature does not contain multiplication. A term $3x$ is just an abbreviation for a term $x + x + x$.

For the semantics I start with considering as the domain the rationals, \mathbb{Q} . As long as coefficients are from the integers, with respect to the satisfiability, validity of a formula the rationals cannot be distinguished from the reals. Restricting the domain from the rationals to the integers, however, results in a difference in satisfiability, validity of a formula, in general. In this case the signature is restricted to integer constants as well.

Definition 6.2.2 (Linear Rational Arithmetic Standard Semantics). The Σ_{LRA} algebra \mathcal{A}_{LRA} is defined by $\text{LA}^{\mathcal{A}_{\text{LRA}}} = \mathbb{Q}$ and all other signature symbols are assigned the standard interpretations over the rationals.

Due to the expressive LA language there is no need for negative literals, because $(\neg <)^{\mathcal{A}_{\text{LRA}}} = (\geq)^{\mathcal{A}_{\text{LRA}}}$, $(\neg >)^{\mathcal{A}_{\text{LRA}}} = (\leq)^{\mathcal{A}_{\text{LRA}}}$, and $(\neg \approx)^{\mathcal{A}_{\text{LRA}}} = (\neq)^{\mathcal{A}_{\text{LRA}}}$.

Note the difference between the above standard semantics over Σ_{LRA} and the free first-order semantics over Σ_{LRA} , Definition 3.2.1. The equation $3 + 4 \approx 5$ has a model in the free first-order semantics, hence it is satisfiable, whereas in the standard model of linear rational arithmetic, Definition 6.2.2, the equation $3 + 4 \approx 5$ is false. In addition, with respect to the standard LRA semantics the definitions of validity, satisfiability coincide with truth and the definition of unsatisfiability coincides with falsehood. This is the result of a single algebra semantics.

6.2.1 Fourier-Motzkin Quantifier Elimination

It is decidable whether a first-order formula over Σ_{LRA} is true or false in the standard LRA semantics. This was first discovered in 1826 by J. Fourier and re-discovered by T. Motzkin in 1936 and is called FM for short. Note that validity of a Σ_{LRA} formula with respect to the standard semantics is undecidable, Exercise ??.

Similar to Congruence Closure, Section 6.1, the starting point of the procedure is a conjunction of atoms without atoms of the form \neq . These will eventually be replaced by a disjunction, i.e., an atom $t \neq s$ is replaced by $t < s \vee t > s$.

Every atom over the variables x, y_1, \dots, y_n can be converted into an equivalent atom $x \circ t[\vec{y}]$ or $0 \circ t[\vec{y}]$, where $\circ \in \{<, >, \leq, \geq, \approx, \neq\}$ and $t[\vec{y}]$ has the form $\sum_i q_i \cdot y_i + q_0$ where $q_i \in \mathbb{Q}$. In other words, a variable x can be either isolated on one side of the atom or eliminated completely. This is the starting point of the FM calculus deciding a conjunction of LA atoms without \neq modulo the isolation of variables and the reduction of ground formulas to \top, \perp .

The calculus operates on a set of atoms N . The normal forms are conjunctions of atoms $s \circ t$ where s, t do not contain any variables. These can be obviously eventually reduced to \top or \perp . The FM calculus consists of two rules:

Substitute $N \uplus \{x \approx t\} \Rightarrow_{\text{FM}} N\{x \mapsto t\}$

provided x does not occur in t

Eliminate $N \uplus \bigcup_i \{x \circ_i^1 t_i\} \uplus \bigcup_j \{x \circ_j^2 s_j\} \Rightarrow_{\text{FM}} N \cup \bigcup_{i,j} \{t_i \circ_{i,j} s_j\}$

provided x does not occur in N nor in the t_i, s_j , $\circ_i^1 \in \{<, \leq\}$, $\circ_j^2 \in \{>, \geq\}$, and $\circ_{i,j} = >$ if $\circ_i^1 = <$ or $\circ_j^2 = >$, and $\circ_{i,j} = \geq$ otherwise

If all variables in N are implicitly existentially quantified, i.e., N stands for $\exists \vec{x}.N$, then the above two rules constitute a sound and complete decision procedure for conjunctions of LA atoms without \neq .

Lemma 6.2.3 (FM Termination on a Conjunction of Atoms). FM terminates on a conjunction of atoms.

Proof. Any rule applications strictly reduces the number of variables. \square

Lemma 6.2.4 (FM Soundness and Completeness on a Conjunction of Atoms). $N \Rightarrow_{\text{FM}}^* \top$ iff $\mathcal{A}_{\text{LRA}} \models \exists \vec{x}.N$. $N \Rightarrow_{\text{FM}}^* \perp$ iff $\mathcal{A}_{\text{LRA}} \not\models \exists \vec{x}.N$.

Proof. \Rightarrow : Assume that $\mathcal{A}_{\text{LRA}}(\beta) \models N$ for some β . Proof by case analysis on the two rules. For rule Substitute obviously $\mathcal{A}_{\text{LRA}}(\beta)(x) = \mathcal{A}_{\text{LRA}}(\beta)(t)$ hence $\mathcal{A}_{\text{LRA}}(\beta) \models N\{x \mapsto t\}$. For rule Eliminate obviously $\mathcal{A}_{\text{LRA}}(\beta)(x) \circ_i^1 \mathcal{A}_{\text{LRA}}(\beta)(t_i)$ and $\mathcal{A}_{\text{LRA}}(\beta)(x) \circ_j^2 \mathcal{A}_{\text{LRA}}(\beta)(s_j)$. A case analysis on \circ_i^1, \circ_j^2 yields $\mathcal{A}_{\text{LRA}}(\beta) \models t_i \circ_{i,j} s_j$ for all i, j .

\Leftarrow : Again by a case analysis on the rules. For rule Substitute if $\mathcal{A}_{\text{LRA}}(\beta) \models N\{x \mapsto t\}$ then $\mathcal{A}_{\text{LRA}}(\beta[x \mapsto \mathcal{A}_{\text{LRA}}(\beta)(t)]) \models N \uplus \{x \approx t\}$. For rule Eliminate if $\mathcal{A}_{\text{LRA}}(\beta) \models N \cup \bigcup_{i,j} \{t_i \circ_{i,j} s_j\}$ then $\mathcal{A}_{\text{LRA}}(\beta[x \mapsto \frac{1}{2}(\min(\bigcup_i \{t_i\}) + \max(\bigcup_j \{s_j\}))]) \models N \uplus \bigcup_i \{x \circ_i^1 t_i\} \uplus \bigcup_j \{x \circ_j^2 s_j\}$. \square

The FM calculus on conjunctions of atoms can be extended to arbitrary closed LRA first-order formulas ϕ . I always assume that different quantifier occurrences in ϕ bind different variables. This can always be obtained by renaming one variable. The first step is to eliminate \top, \perp from ϕ and to transform ϕ in negation normal form, see Section 3.9. The resulting formula only contains the operators $\forall, \exists, \wedge, \vee, \neg$, where all negation symbols occur in front of atoms. Then the following rule can be used to remove the negation symbols as well:

$$\mathbf{ElimNeg} \quad \chi[\neg s \circ^1 t]_p \Rightarrow_{\text{FM}} \chi[s \circ^2 t]_p$$

where the pairs (\circ_1, \circ_2) are given by pairs $(<, \geq), (\leq, >), (\approx, \neq)$ and their symmetric variants

The above two FM rules on conjunctions cannot cope with atoms $s \neq t$, so they are eliminated as well:

$$\mathbf{Elim}\neq \quad \chi[s \neq t]_p \Rightarrow_{\text{FM}} \chi[s < t \vee s > t]_p$$

The next step is to compute a *Prenex Normal Form*, a formula $\{\exists, \forall\}x_1 \dots \{\exists, \forall\}x_n.\phi$ where ϕ does not contain any quantifiers. This can be done by simply applying the mini-scoping rules, see Section 3.9, in the opposite direction:

Prenex1 $\chi[(\forall x.\psi_1) \circ \psi_2]_p \Rightarrow_{\text{FM}} \chi[\forall x.(\psi_1 \circ \psi_2)]_p$
 provided $\circ \in \{\wedge, \vee\}$, $x \notin \text{fvars}(\psi_2)$

Prenex2 $\chi[(\exists x.\psi_1) \circ \psi_2]_p \Rightarrow_{\text{FM}} \chi[\exists x.(\psi_1 \circ \psi_2)]_p$
 provided $\circ \in \{\wedge, \vee\}$, $x \notin \text{fvars}(\psi_2)$

Prenex3 $\chi[(\forall x.\psi_1) \wedge (\forall y.\psi_2)]_p \Rightarrow_{\text{FM}} \chi[\forall x.(\psi_1 \wedge \psi_2\{y \mapsto x\})]_p$

Prenex4 $\chi[(\exists x.\psi_1) \vee (\exists y.\psi_2)]_p \Rightarrow_{\text{FM}} \chi[\exists x.(\psi_1 \vee \psi_2\{y \mapsto x\})]_p$

where Prenex3 and Prenex4 are preferred over Prenex1 and Prenex2. Finally, for the resulting formula $\{\exists, \forall\}x_1 \dots \{\exists, \forall\}x_n.\phi$ in prenex normal form the FM algorithm computes a DNF of ϕ by exhaustively applying the rule PushConj, Section 2.5.2. The result is a formula $\{\exists, \forall\}x_1 \dots \{\exists, \forall\}x_n.\phi$ where ϕ is a DNF of atoms without containing an atom of the form $s \not\approx t$. All the above formulas transformations are equivalence preserving. Therefore, to each conjunct of ϕ the above two FM rules decide the conjunct, if all variables are existentially quantified. This is the final obstacle in order to obtain the FM decision procedure for arbitrary formulas.

It is solved by considering the quantifiers iteratively in an innermost way. For the formula $\{\exists, \forall\}x_1 \dots \{\exists, \forall\}x_n.\phi$ always the innermost quantifier $\{\exists, \forall\}x_n$ is considered. If it is an existential quantifier, $\exists x_n$, then the FM rules Substitute, Eliminate are applied to the variable x_n for each conjunct C_i of $\phi = C_1 \vee \dots \vee C_n$. The result is a formula $\{\exists, \forall\}x_1 \dots \{\exists, \forall\}x_{n-1}.(C'_1 \vee \dots \vee C'_n)$ which is again in prenex DNF. Furthermore, by Lemma 6.2.4 it is equivalent to $\{\exists, \forall\}x_1 \dots \{\exists, \forall\}x_n.\phi$. If the innermost quantifier is a universal quantifier $\forall x_n$, then the formula is replaced by $\{\exists, \forall\}x_1 \dots \{\exists, \forall\}x_{n-1}.\neg \exists x_n.\neg \phi$ and the above steps for negation normal form and DNF are repeated for $\neg \phi$ resulting in an equivalent formula $\{\exists, \forall\}x_1 \dots \{\exists, \forall\}x_{n-1}.\neg \exists x_n.\phi'$ where ϕ' is in DNF and does not contain negation symbols nor atoms $s \not\approx t$. Then the FM rules Substitute, Eliminate are applied to the variable x_n for each conjunct C_i of $\phi' = C_1 \vee \dots \vee C_n$. The result is an equivalent formula $\{\exists, \forall\}x_1 \dots \{\exists, \forall\}x_{n-1}.\neg(C'_1 \vee \dots \vee C'_n)$. Finally, the above steps for negation normal form and DNF are repeated for $\neg(C'_1 \vee \dots \vee C'_n)$ resulting in an equivalent formula $\{\exists, \forall\}x_1 \dots \{\exists, \forall\}x_{n-1}.\phi''$ where ϕ'' is in DNF and does not contain negation symbols nor atoms $s \not\approx t$. This completes for FM decision procedure for LRA formulas.

Every LRA formula can be reduced to \top or \perp via the FM decision procedure. Therefore LRA is called a *complete* theory, i.e., every closed formula over the signature of LRA is either true or false.

LA formulas over the rationals and over the reals are indistinguishable by first-order formulas over the signature of LRA. These properties do not hold for extended signatures, e.g., then additional free symbols are introduced. Furthermore, FM is no decision procedures over the integers, even if the LA syntax is restricted to integer constants.

The complexity of the FM calculus depends mostly on the quantifier alternations in $\{\exists, \forall\}x_1 \dots \{\exists, \forall\}x_n.\phi$. In case an existential quantifier \exists is eliminated, the formula size grows worst-case quadratically, therefore $O(n^2)$ runtime. For m quantifiers $\exists \dots \exists$: a naive implementation needs worst-case $O(n^{2^m})$ runtime. It is not known whether an optimized implementation with simply exponential runtime is possible. If there are m quantifier alternations $\exists\forall\exists\forall \dots \exists\forall$, a CNF to DNF conversion is required after each step. Each conversion has a worst-case exponential run time, see Section 2.5. Therefore, the overall procedure has a worst-case non-elementary runtime.

I There are meanwhile more efficient decision procedures for the theory LRA known, e.g., see Section 6.2.3. There are problems occurring in practice where the elimination of a variable via FM results in an only linear increase in size. In such cases FM is still valuable. Many state-of-the-art LRA procedures actually calculate the size of the formula after eliminating a variable via FM and redundancy elimination and decide on this basis whether FM is applied or not.