

3.1.2 Definition (Term)

Given a signature $\Sigma = (\mathcal{S}, \Omega, \Pi)$, a sort $S \in \mathcal{S}$ and a variable set \mathcal{X} , the set $T_S(\Sigma, \mathcal{X})$ of all *terms* of sort S is recursively defined by (i) $x_S \in T_S(\Sigma, \mathcal{X})$ if $x_S \in \mathcal{X}$, (ii) $f(t_1, \dots, t_n) \in T_S(\Sigma, \mathcal{X})$ if $f \in \Omega$ and $f : S_1 \times \dots \times S_n \rightarrow S$ and $t_i \in T_{S_i}(\Sigma, \mathcal{X})$ for every $i \in \{1, \dots, n\}$.

The sort of a term t is denoted by $\text{sort}(t)$, i.e., if $t \in T_S(\Sigma, \mathcal{X})$ then $\text{sort}(t) = S$. A term not containing a variable is called *ground*.

For the sake of simplicity it is often written: $T(\Sigma, \mathcal{X})$ for $\bigcup_{S \in \mathcal{S}} T_S(\Sigma, \mathcal{X})$, the set of all terms, $T_S(\Sigma)$ for the set of all ground terms of sort $S \in \mathcal{S}$, and $T(\Sigma)$ for $\bigcup_{S \in \mathcal{S}} T_S(\Sigma)$, the set of all ground terms over Σ .

Note that the sets $T_S(\Sigma)$ are all non-empty, because there is at least one constant for each sort S in Σ . The sets $T_S(\Sigma, \mathcal{X})$ include infinitely many variables of sort S .



3.1.3 Definition (Equation, Atom, Literal)

If $s, t \in T_S(\Sigma, \mathcal{X})$ then $s \approx t$ is an *equation* over the signature Σ . Any equation is an *atom* (also called *atomic formula*) as well as every $P(t_1, \dots, t_n)$ where $t_i \in T_{S_i}(\Sigma, \mathcal{X})$ for every $i \in \{1, \dots, n\}$ and $P \in \Pi$, $\text{arity}(P) = n$, $P \subseteq S_1 \times \dots \times S_n$.

An atom or its negation of an atom is called a *literal*.

Definition (Formulas)

The set $FOL(\Sigma, \mathcal{X})$ of *many-sorted first-order formulas with equality* over the signature Σ is defined as follows for formulas $\phi, \psi \in F_{\Sigma}(\mathcal{X})$ and a variable $x \in \mathcal{X}$:

$FOL(\Sigma, \mathcal{X})$	Comment
\perp	false
\top	true
$P(t_1, \dots, t_n), s \approx t$	atom
$(\neg\phi)$	negation
$(\phi \circ \psi)$	$\circ \in \{\wedge, \vee, \rightarrow, \leftrightarrow\}$
$\forall x.\phi$	universal quantification
$\exists x.\phi$	existential quantification

?? Definition (Positions)

The set of positions of a term, formula is inductively defined by:

$$\text{pos}(x) := \{\epsilon\} \text{ if } x \in \mathcal{X}$$

$$\text{pos}(\phi) := \{\epsilon\} \text{ if } \phi \in \{\top, \perp\}$$

$$\text{pos}(\neg\phi) := \{\epsilon\} \cup \{1p \mid p \in \text{pos}(\phi)\}$$

$$\text{pos}(\phi \circ \psi) := \{\epsilon\} \cup \{1p \mid p \in \text{pos}(\phi)\} \cup \{2p \mid p \in \text{pos}(\psi)\}$$

$$\text{pos}(s \approx t) := \{\epsilon\} \cup \{1p \mid p \in \text{pos}(s)\} \cup \{2p \mid p \in \text{pos}(t)\}$$

$$\text{pos}(f(t_1, \dots, t_n)) := \{\epsilon\} \cup \bigcup_{i=1}^n \{ip \mid p \in \text{pos}(t_i)\}$$

$$\text{pos}(P(t_1, \dots, t_n)) := \{\epsilon\} \cup \bigcup_{i=1}^n \{ip \mid p \in \text{pos}(t_i)\}$$

$$\text{pos}(\forall x.\phi) := \{\epsilon\} \cup \{1p \mid p \in \text{pos}(\phi)\}$$

$$\text{pos}(\exists x.\phi) := \{\epsilon\} \cup \{1p \mid p \in \text{pos}(\phi)\}$$

where $\circ \in \{\wedge, \vee, \rightarrow, \leftrightarrow\}$ and $t_i \in T(\Sigma, \mathcal{X})$ for all $i \in \{1, \dots, n\}$.

An term t (formula ϕ) is said to *contain* another term s (formula ψ) if $t|_p = s$ ($\phi|_p = \psi$). It is called a *strict subexpression* if $p \neq \epsilon$. The term t (formula ϕ) is called an *immediate subexpression* of s (formula ψ) if $|p| = 1$. For terms a subexpression is called a *subterm* and for formulas a *subformula*, respectively.

The *size* of a term t (formula ϕ), written $|t|$ ($|\phi|$), is the cardinality of $\text{pos}(t)$, i.e., $|t| := |\text{pos}(t)|$ ($|\phi| := |\text{pos}(\phi)|$). The *depth* of a term, formula is the maximal length of a position in the term, formula: $\text{depth}(t) := \max\{|p| \mid p \in \text{pos}(t)\}$ ($\text{depth}(\phi) := \max\{|p| \mid p \in \text{pos}(\phi)\}$).



The set of *all* variables occurring in a term t (formula ϕ) is denoted by $\text{vars}(t)$ ($\text{vars}(\phi)$) and formally defined as

$$\text{vars}(t) := \{x \in \mathcal{X} \mid x = t|_p, p \in \text{pos}(t)\}$$

$$(\text{vars}(\phi) := \{x \in \mathcal{X} \mid x = \phi|_p, p \in \text{pos}(\phi)\}).$$

A term t (formula ϕ) is *ground* if $\text{vars}(t) = \emptyset$ ($\text{vars}(\phi) = \emptyset$). Note that $\text{vars}(\forall x.a \approx b) = \emptyset$ where a, b are constants. This is justified by the fact that the formula does not depend on the quantifier, see the semantics below. The set of *free* variables of a formula ϕ (term t) is given by $\text{fvvars}(\phi, \emptyset)$ ($\text{fvvars}(t, \emptyset)$) and recursively defined by $\text{fvvars}(\psi_1 \circ \psi_2, B) := \text{fvvars}(\psi_1, B) \cup \text{fvvars}(\psi_2, B)$ where $\circ \in \{\wedge, \vee, \rightarrow, \leftrightarrow\}$, $\text{fvvars}(\forall x.\psi, B) := \text{fvvars}(\psi, B \cup \{x\})$, $\text{fvvars}(\exists x.\psi, B) := \text{fvvars}(\psi, B \cup \{x\})$, $\text{fvvars}(\neg\psi, B) := \text{fvvars}(\psi, B)$, $\text{fvvars}(L, B) := \text{vars}(L) \setminus B$ ($\text{fvvars}(t, B) := \text{vars}(t) \setminus B$). For $\text{fvvars}(\phi, \emptyset)$ I also write $\text{fvvars}(\phi)$.

In $\forall x.\phi$ ($\exists x.\phi$) the formula ϕ is called the *scope* of the quantifier. An occurrence q of a variable x in a formula ϕ ($\phi|_q = x$) is called *bound* if there is some $p < q$ with $\phi|_p = \forall x.\phi'$ or $\phi|_p = \exists x.\phi'$. Any other occurrence of a variable is called *free*.

A formula not containing a free occurrence of a variable is called *closed*. If $\{x_1, \dots, x_n\}$ are the variables freely occurring in a formula ϕ then $\forall x_1, \dots, x_n.\phi$ and $\exists x_1, \dots, x_n.\phi$ (abbreviations for $\forall x_1.\forall x_2 \dots \forall x_n.\phi$, $\exists x_1.\forall x_2 \dots \forall x_n.\phi$, respectively) are the *universal* and the *existential closure* of ϕ , respectively.



3.1.7 Definition (Polarity)

The *polarity* of a subformula $\psi = \phi|_p$ at position p is $pol(\phi, p)$ where pol is recursively defined by

$$\begin{aligned}
 pol(\phi, \epsilon) &:= 1 \\
 pol(\neg\phi, 1p) &:= -pol(\phi, p) \\
 pol(\phi_1 \circ \phi_2, ip) &:= pol(\phi_i, p) \text{ if } \circ \in \{\wedge, \vee\} \\
 pol(\phi_1 \rightarrow \phi_2, 1p) &:= -pol(\phi_1, p) \\
 pol(\phi_1 \rightarrow \phi_2, 2p) &:= pol(\phi_2, p) \\
 pol(\phi_1 \leftrightarrow \phi_2, ip) &:= 0 \\
 pol(P(t_1, \dots, t_n), p) &:= 1 \\
 pol(t \approx s, p) &:= 1 \\
 pol(\forall x.\phi, 1p) &:= pol(\phi, p) \\
 pol(\exists x.\phi, 1p) &:= pol(\phi, p)
 \end{aligned}$$

Semantics

3.2.1 Definition (Σ -algebra)

Let $\Sigma = (\mathcal{S}, \Omega, \Pi)$ be a signature with set of sorts \mathcal{S} , operator set Ω and predicate set Π . A Σ -algebra \mathcal{A} , also called Σ -interpretation, is a mapping that assigns (i) a non-empty carrier set $S^{\mathcal{A}}$ to every sort $S \in \mathcal{S}$, so that $(S_1)^{\mathcal{A}} \cap (S_2)^{\mathcal{A}} = \emptyset$ for any distinct sorts $S_1, S_2 \in \mathcal{S}$, (ii) a total function $f^{\mathcal{A}} : (S_1)^{\mathcal{A}} \times \dots \times (S_n)^{\mathcal{A}} \rightarrow (S)^{\mathcal{A}}$ to every operator $f \in \Omega$, $\text{arity}(f) = n$ where $f : S_1 \times \dots \times S_n \rightarrow S$, (iii) a relation $P^{\mathcal{A}} \subseteq ((S_1)^{\mathcal{A}} \times \dots \times (S_m)^{\mathcal{A}})$ to every predicate symbol $P \in \Pi$, $\text{arity}(P) = m$. (iv) the equality relation becomes $\approx^{\mathcal{A}} = \{(e, e) \mid e \in \mathcal{U}^{\mathcal{A}}\}$ where the set $\mathcal{U}^{\mathcal{A}} := \bigcup_{S \in \mathcal{S}} (S)^{\mathcal{A}}$ is called the *universe* of \mathcal{A} .

A (variable) *assignment*, also called a *valuation* for an algebra \mathcal{A} is a function $\beta : \mathcal{X} \rightarrow \mathcal{U}_{\mathcal{A}}$ so that $\beta(x) \in S_{\mathcal{A}}$ for every variable $x \in \mathcal{X}$, where $S = \text{sort}(x)$. A *modification* $\beta[x \mapsto e]$ of an assignment β at a variable $x \in \mathcal{X}$, where $e \in S_{\mathcal{A}}$ and $S = \text{sort}(x)$, is the assignment defined as follows:

$$\beta[x \mapsto e](y) = \begin{cases} e & \text{if } x = y \\ \beta(y) & \text{otherwise.} \end{cases}$$



The homomorphic extension $\mathcal{A}(\beta)$ of β onto terms is a mapping $T(\Sigma, \mathcal{X}) \rightarrow \mathcal{U}_{\mathcal{A}}$ defined as (i) $\mathcal{A}(\beta)(x) = \beta(x)$, where $x \in \mathcal{X}$ and (ii) $\mathcal{A}(\beta)(f(t_1, \dots, t_n)) = f_{\mathcal{A}}(\mathcal{A}(\beta)(t_1), \dots, \mathcal{A}(\beta)(t_n))$, where $f \in \Omega$, $\text{arity}(f) = n$.

Given a term $t \in T(\Sigma, \mathcal{X})$, the value $\mathcal{A}(\beta)(t)$ is called the *interpretation* of t under \mathcal{A} and β . If the term t is ground, the value $\mathcal{A}(\beta)(t)$ does not depend on a particular choice of β , for which reason the interpretation of t under \mathcal{A} is denoted by $\mathcal{A}(t)$. An algebra \mathcal{A} is called *term-generated*, if every element e of the universe $\mathcal{U}_{\mathcal{A}}$ of \mathcal{A} is the image of some ground term t , i.e., $\mathcal{A}(t) = e$.

3.2.2 Definition (Semantics)

An algebra \mathcal{A} and an assignment β are extended to formulas $\phi \in \text{FOL}(\Sigma, \mathcal{X})$ by

$$\mathcal{A}(\beta)(\perp) := 0 \quad \mathcal{A}(\beta)(\top) := 1$$

$$\mathcal{A}(\beta)(s \approx t) := 1 \text{ if } \mathcal{A}(\beta)(s) = \mathcal{A}(\beta)(t) \text{ else } 0$$

$$\mathcal{A}(\beta)(P(t_1, \dots, t_n)) := 1 \text{ if } (\mathcal{A}(\beta)(t_1), \dots, \mathcal{A}(\beta)(t_n)) \in P_{\mathcal{A}} \text{ else } 0$$

$$\mathcal{A}(\beta)(\neg\phi) := 1 - \mathcal{A}(\beta)(\phi)$$

$$\mathcal{A}(\beta)(\phi \wedge \psi) := \min(\{\mathcal{A}(\beta)(\phi), \mathcal{A}(\beta)(\psi)\})$$

$$\mathcal{A}(\beta)(\phi \vee \psi) := \max(\{\mathcal{A}(\beta)(\phi), \mathcal{A}(\beta)(\psi)\})$$

$$\mathcal{A}(\beta)(\phi \rightarrow \psi) := \max(\{(1 - \mathcal{A}(\beta)(\phi)), \mathcal{A}(\beta)(\psi)\})$$

$$\mathcal{A}(\beta)(\phi \leftrightarrow \psi) := \text{if } \mathcal{A}(\beta)(\phi) = \mathcal{A}(\beta)(\psi) \text{ then } 1 \text{ else } 0$$

$$\mathcal{A}(\beta)(\exists x_S.\phi) := 1 \text{ if } \mathcal{A}(\beta[x \mapsto e])(\phi) = 1$$

for some $e \in S_{\mathcal{A}}$ and 0 otherwise

$$\mathcal{A}(\beta)(\forall x_S.\phi) := 1 \text{ if } \mathcal{A}(\beta[x \mapsto e])(\phi) = 1$$

for all $e \in S_{\mathcal{A}}$ and 0 otherwise



A formula ϕ is called *satisfiable by \mathcal{A} under β* (or *valid in \mathcal{A} under β*) if $\mathcal{A}, \beta \models \phi$; in this case, ϕ is also called *consistent*;

satisfiable by \mathcal{A} if $\mathcal{A}, \beta \models \phi$ for some assignment β ;

satisfiable if $\mathcal{A}, \beta \models \phi$ for some algebra \mathcal{A} and some assignment β ;

valid in \mathcal{A} , written $\mathcal{A} \models \phi$, if $\mathcal{A}, \beta \models \phi$ for any assignment β ; in this case, \mathcal{A} is called a *model* of ϕ ;

valid, written $\models \phi$, if $\mathcal{A}, \beta \models \phi$ for any algebra \mathcal{A} and any assignment β ; in this case, ϕ is also called a *tautology*;

unsatisfiable if $\mathcal{A}, \beta \not\models \phi$ for any algebra \mathcal{A} and any assignment β ; in this case ϕ is also called *inconsistent*.

